arXiv:1702.03819v2 [math.GT] 14 Feb 2017

p-ADIC MAHLER MEASURE AND $\mathbb{Z}$-COVERS OF LINKS

JUN UEKI

Abstract. Let p be a prime number. We develop a theory of p-adic Mahler measure of polynomials and apply it to the study of $\mathbb{Z}$-covers of rational homology 3-spheres branched over links. We obtain a p-adic analogue of the asymptotic formula of the torsion homology growth and a balance formula among the leading coefficient of the Alexander polynomial, the p-adic entropy, and the Iwasawa μ_p-invariant. We also apply the purely p-adic theory of Besser–Deninger to $\mathbb{Z}$-covers of links. In addition, we study the entropies of profinite cyclic covers of links. We examine various examples throughout the paper.

Contents

1. Introduction 2
2. p-adic Mahler measure of polynomials 3
2.1. Euclidean Mahler measure 3
2.2. p-adic Mahler measure M_p 4
2.3. Proofs 6
2.4. Alternative modification of the integral path 7
3. p-adic Mahler measure and $\mathbb{Z}$-covers of links 8
3.1. $\mathbb{Z}$-covers and M_p 8
3.2. Iwasawa μ_p-invariant and M_p 9
3.3. p-adic entropy h_p and M_p 10
3.4. h_p, μ_p, and the leading coefficient a_0 11
4. Purely p-adic notions 12
4.1. Besser–Deninger m_p 12
4.2. Deninger's $\hbar_p$ 13
5. Profinite cyclic covers 14
5.1. $\mathbb{Z}_p$-covers 14
5.2. $\widehat{\mathbb{Z}}$-covers 15
Acknowledgment 16
References 16

Date: February 15, 2017.
2010 *Mathematics Subject Classification.* Primary 37B40, 57M12, 11R06; Secondary 11R23.
Key words and phrases. link, branched covering, Mahler measure, entropy, p-adic.

1. Introduction

It is well known that the asymptotic behavior of the torsion homology growth in a $\mathbb{Z}$-cover of S^3 branched over a link is described by the Euclidean Mahler measure of the reduced Alexander polynomial of the branch link.

The Euclidean Mahler measure $\mathrm{M}(f(t))$ of a Laurent polynomial $0 \neq f(t) \in \mathbb{C}[t^{\pm 1}]$ is defined by $\log \mathrm{M}(f(t)) = \dfrac{1}{2\pi\sqrt{-1}} \displaystyle\int_{|z|=1} \log|f(z)| \frac{dz}{z}$. If $f(t) = a\prod_i (t-\alpha_i)$, then we have Jensen's formula $\mathrm{M}(f(t)) = |a| \prod_i \max\{|\alpha_i|, 1\}$ ([EW99]).

If H is a finite group, then we denote its order by $|H|$. If H is an infinite group, then we put $|H| = 0$. For two polynomials $f(t)$ and $g(t)$ in $\mathbb{Z}[t]$, we denote their resultant by $R(f(t), g(t))$. Now let M be a rational homology 3-sphere ($\mathbb{Q}\mathrm{HS}^3$) and let L be a link in M. Let $h_\infty : X_\infty \to X := M - L$ be a $\mathbb{Z}$-cover over the exterior with a generator t of the deck transformation group and let $\{h_n : M_n \to M\}_n$ denote the system of branched $\mathbb{Z}/n\mathbb{Z}$-covers over (M, L) obtained from subcovers of h_∞ by the Fox completion. The Alexander module $H_1(X_\infty)$ is a finitely generated $\mathbb{Z}[t^{\mathbb{Z}}]$-module. We take a generator $A_L(t)$ of the maximum principal ideal containing the Fitting ideal $\mathrm{Fitt}_{\mathbb{Z}[t^{\mathbb{Z}}]} H_1(X_\infty)$ of $H_1(X_\infty)$ over $\mathbb{Z}[t^{\mathbb{Z}}]$ with $A_L(t) \in \mathbb{Z}[t] \subset \mathbb{Z}[t^{\mathbb{Z}}]$ and call it *an/the Alexander polynomial of* h_∞. Put $\nu_n(t) := (t^n - 1)/(t - 1)$. Then by a generalization of Fox's formula (Proposition 3.2, e.g.[KM08]), there exists some finite set $\mathcal{C} \subset \mathbb{N}$ such that for any $n \gg 0$ the resultant $R(A_L(t), \nu_n(t))$ satisfies $|H_1(M_n)| = |R(A_L(t), \nu_n(t))| \times C$ for some $C \in \mathcal{C}$. By a similar argument to [GAS91], we obtain an asymptotic formula of the homology growth (Theorem 3.3): $\lim\limits_{n\in\mathbb{N};\mathrm{value}\neq 0} |H_1(M_n)|^{1/n} = \mathrm{M}(A_L(t))$.

For a prime number p, let $|\cdot|_p$ denote the p-adic norm (absolute value) normalized by $|p|_p = p^{-1}$. If H is a group, then we write $||H||_p = |(|H|)|_p$. For an inverse system $\{M_{p^r} \to M\}_r$ of branched $\mathbb{Z}/p^r\mathbb{Z}$-covers over (M, L), a p-adic refinement of the asymptotic formula called *the Iwasawa type formula* (Proposition 3.6) is known: If M_{p^r}'s are $\mathbb{Q}\mathrm{HS}^3$'s, then there exist some $\lambda_p, \mu_p \in \mathbb{N}, \nu_p \in \mathbb{Z}$ called the Iwasawa invariants such that for any $r \gg 0$ the equation $||H_1(M_{p^r})||_p^{-1} = p^{\lambda_p r + \mu_p p^r + \nu_p}$ holds.

The main purpose of this paper is to investigate the relation between the Mahler measure and the Iwasawa invariants.

For a prime number p, let $\mathbb{C}_p = \widehat{\overline{\mathbb{Q}}}_p$ denote the p-adic completion of an algebraic closure of the p-adic number field $\mathbb{Q}_p$, and fix an immersion $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ of an algebraic closure of $\mathbb{Q}$. (We regard $p = \infty$ for Euclidean objects.)

We define the p-adic Mahler measure $\mathrm{M}_p(f(z))$ of a Laurent polynomial $f(t) \in \mathbb{C}_p[t^{\pm 1}]$ ($\neq 0$) by imitating the Shnirel'man integral ([Shn38]) as follows: We first suppose that $f(t)$ does not vanish on the p-adic unit circle $|z|_p = 1$ and put

$$\log \mathrm{M}_p(f(z)) := \lim_{n\in\mathbb{N};\gcd(n,p)=1} \frac{1}{n} \sum_{\zeta^n=1} \log|f(\zeta)|_p,$$

where the limit is taken with respect to the Euclidean topology.

If $f(t) = \sum_{0\leq i\leq l} a_i t^{d-i} = a_0 t^{d-l} \prod_i (t - \alpha_i)$, then an analogue $\mathrm{M}_p(f(t)) = |a_0|_p \prod_i \max\{|\alpha_i|_p, 1\}$ of Jensen's formula holds (Theorem 2.3). In addition, the equation $\mathrm{M}_p(f(t)) = \max\{|a_i|_p\}_i$ holds (Proposition 2.7), that is, $\mathrm{M}_p(f(t))$ coincides with the Gauss norm of $f(t)$.

We extend this notion with Jensen's formula to the case with roots on $|z|_p = 1$ and remove the condition $\gcd(n,p) = 1$ on the limit (Theorem 2.5). Then for a $\mathbb{Z}$-cover over (M,L) with its polynomial $A_L(t)$, we obtain a p-adic analogue $\lim_{n\in\mathbb{N};\text{value}\neq 0} ||H_1(M_n)||_p^{1/n} = \mathrm{M}_p(A_L(t))$ of the asymptotic formula (Theorem 3.3). In addition, the formula $\log \mathrm{M}_p(A_L(t)) = -\mu_p \log p$ of the Iwasawa invariant follows (Proposition 3.7).

Now we consider a $\mathbb{Z}$-cover over (S^3, L) with $A_L(t) = \sum_{0\le i\le l} a_i t^{d-i} = a_0 t^{d-l} \prod_i (t-\alpha_i)$ over $\overline{\mathbb{Q}}$. The relation between the Mahler measure and the topological entropy is well known. It is also known that a_0 has a topological information of L. Noguchi studied the relation among them from a viewpoint of p-adic topology ([Nog07]). We generalize his work for links.

We consider the meridian action on $H_1(X_\infty, \mathbb{Q})$ and its Pontryagin dual. Then by [GBV15] and [LW88], the algebraic/topological entropy is given by $h = \sum_{p\le\infty} h_p = \sum_{p\le\infty} \log \mathrm{M}_p(A_L(t))$ with use of the p-adic entropies $h_p = \log \mathrm{M}_p(A_L(t)/a_0) = \sum_{|\alpha_i|_p>1} \log |\alpha_i|_p$ (Proposition 3.11). We see a balance formula among a_0, h_p, and the Iwasawa μ_p-invariant (Proposition 3.13):

$$-\log |a_0|_p = h_p + \mu_p \log p.$$

By the product formula $\prod_{p\le\infty} |a_0|_p = 1$, a generalization of [Nog07, Corollary 4] follows: $\log |a_0| = \sum_{p<\infty} \sum_{|\alpha_i|_p>1} \log |\alpha_i|_p + \sum_{p<\infty} \mu_p \log p$ (Corollary 3.15).

Besser and Deninger ([BD99]) defined the *purely* p-adic log Mahler measure m_p, which is different from ours, with use of the p-adic logarithm and the Shnirel'man integral, and proved Jensen's formula for it. In addition, Deninger ([Den09]) introduced the notion of the *purely* p-adic entropy $\hbar_p$ by using the numbers of fixed points and the p-adic logarithm, and proved the relation between m_p and $\hbar_p$. In Section 4, we apply their theory to $\mathbb{Z}$-covers of S^3 branched over links.

Finally, we study profinite cyclic covers of S^3 branched over links. We consider the p-adic integer ring $\mathbb{Z}_p = \varprojlim_n \mathbb{Z}/p^n\mathbb{Z}$ and the profinite integer ring $\widehat{\mathbb{Z}} = \varprojlim_n \mathbb{Z}/n\mathbb{Z}$. We investigate the entropies of the meridian actions on the Iwasawa modules $\mathcal{H}_p$ of $\mathbb{Z}_p$-covers and those on the completed Alexander modules $\mathcal{H}$ of $\widehat{\mathbb{Z}}$-covers for the cases with $A_L(t) \in \mathbb{Z}[t]$.

In Sections 3–5, we compute various examples obtained from links in the Rolfsen table ([Rol76]).

2. p-adic Mahler measure of polynomials

2.1. **Euclidean Mahler measure.** We consider the complex function on $\mathbb{C}^*$ given by a Laurent polynomial $0 \neq f(t) \in \mathbb{C}[t^{\pm 1}]$. If $f(t) = \sum_{0\le i\le l} a_i t^{d-i} = a_0 t^{d-l} \prod_i (t-\alpha_i)$ with $a_0 a_l \neq 0$ and it has no root on the unit circle $|z| = 1$, then its Euclidean Mahler measure $\mathrm{M}(f(t))$ is defined by

$$\log \mathrm{M}(f(t)) = \frac{1}{2\pi\sqrt{-1}} \int_{|z|=1} \log |f(z)| \frac{dz}{z}.$$

It satisfies Jensen's formula

$$\begin{aligned}\log \mathrm{M}(f(t)) &= \log|a_l| - \sum_{0<|\alpha_i|<1} \log|\alpha_i| \\ &= \log|a_0| + \sum_{1<|\alpha_i|} \log|\alpha_i|.\end{aligned}$$

It is generalized to the cases with roots on $|z|=1$. We define $\mathrm{M}(f(t))$ as the limit of the integrals along paths avoiding roots on $|z|=1$ [EW99]. Indeed, suppose $f(\alpha)=0, |\alpha|=1$. Then for each simple closed curve γ around α with finite length $|\gamma|$, the integration of a bounded continuous function $\log|f(z)|/z$ along γ is bounded by $|\gamma| \times$ (constant). Thus we have $\lim_{|\gamma|\to 0} \int_\gamma \log|f(z)| \frac{dz}{z} = 0$.

For polynomials $f(t) = a\prod_i (t-\alpha_i)$, $g(t) = b\prod_j (t-\beta_j) \in \mathbb{Z}[t]$, their resultant is defined as

$$R(f(t), g(t)) := a^{\deg(g)} b^{\deg(f)} \prod_{i,j} (\alpha_i - \beta_j).$$

It coincides with the determinant of the Sylvester matrix, whose elements are given by their coefficients, and hence satisfies $R(f(t),g(t)) \in \mathbb{Z}$. We have $R(f(t),g(t))=0$ if and only if they gave no common zero. We have

$$\begin{aligned}R(f(t),g(t)) &= a^{\deg(g)} \prod_i g(\alpha_i) = b^{\deg(f)} \prod_j f(\beta_j), \\ R(f(t),g(t)) &= (-1)^{\deg(fg)} R(g(t),f(t)), \\ R(f(t),g(t)h(t)) &= R(f(t),g(t)) R(f(t),h(t))\end{aligned}$$

for each $h(t) \in \mathbb{Z}[t]$. If $f(t)=0$, then we put $R(0,g(t))=0$. If $g(t) \neq 0$, then we put $R(1,g(t))=1$ ([Web79]).

By an argument in [GAS91], we have an asymptotic formula of the resultants:

Proposition 2.1. Let $0 \neq f(t) \in \mathbb{Z}[t]$ and $\nu_n(t) = t^n - 1 = \sum_{0\le i<n} t^i$. Then

$$\lim_{n\in\mathbb{N};\mathrm{value}\neq 0} |R(f(t),\nu_n(t))|^{1/n} = \mathrm{M}(f(t)).$$

In order to prove this, we need to show $\lim_{n\in\mathbb{N};\mathrm{value}\neq 0} |\alpha^n - 1|^{1/n} = 1$ for a root α with $|\alpha|=1$. In [GAS91], the notion of $\arg\alpha$ was used in obtaining the estimation $|\alpha^n - 1| > C\exp(-(\log n)^6)$ for a constant C.

2.2. **p-adic Mahler measure M_p.** For a prime number p, let $|\cdot|_p$ denote the p-adic norm normalized by $|p|_p = 1/p$, and let $\mathbb{Q}_p$ denote the p-adic numbers, that is, the p-adic completion of $\mathbb{Q}$. Let $\mathbb{C}_p = \widehat{\overline{\mathbb{Q}}}_p$ denote the p-adic completion of an algebraic closure of $\mathbb{Q}_p$, and fix an immersion $\overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ of an algebraic closure of $\mathbb{Q}$. For a continuous function $F : \{z \in \mathbb{C}_p \mid |z|_p = 1\} \to \mathbb{C}_p$ with no zero on $|z|_p = 1$, the Shnirel'man integral is defined by

$$\int_{|z|_p=1} F(z)\frac{dz}{z} := \lim_{n\in\mathbb{N};\gcd(n,p)=1} \frac{1}{n} \sum_{\zeta^n=1} F(\zeta),$$

where the limit is taken with respect to the p-adic topology ([Shn38]).

We *imitate* this notion to define the p-adic Mahler measure as follows:

Definition 2.2. Let $f : \mathbb{C}_p^* \to \mathbb{C}_p$ be the continuous map defined by a Laurent polynomial $0 \neq f(t) \in \mathbb{C}_p[t^{\pm 1}]$. If $f(t)$ has no root on $|z|_p = 1$, then we define its *p-adic Mahler measure* $\mathrm{M}_p(f(t))$ by

$$\log \mathrm{M}_p(f(t)) = \lim_{n \in \mathbb{N}; \gcd(n,p)=1} \frac{1}{n} \sum_{\zeta^n = 1} \log |f(\zeta)|_p,$$

where the limit is taken with respect to the Euclidean topology.

Then we have an analogue of Jensen's formula:

Theorem 2.3 (Jensen's formula)**.** If $f(t) = \sum_{0 \leq i \leq l} a_i t^{d-i} = a_0 t^{d-l} \prod_i (t - \alpha_i) \in \mathbb{C}_p[t^{\pm 1}]$ with $a_0 a_l \neq 0$, then

$$\begin{aligned} \log \mathrm{M}_p(f(t)) &= \log |a_l|_p - \sum_{0 < |\alpha_i|_p < 1} \log |\alpha_i|_p \\ &= \log |a_0|_p + \sum_{1 < |\alpha_i|_p} \log |\alpha_i|_p. \end{aligned}$$

Remark 2.4. Besser and Deninger defined the *purely* p-adic Mahler measure with use of the Shnirel'man integral and the p-adic log, which is different from ours, and proved an analogue of Jensen's formula for it ([BD99]). We recall it in Section 4.

For our $\mathrm{M}_p(f(t))$, we can remove the condition $\gcd(n, p) = 1$ on the limit. In addition, we can extend this notion for $f(t)$ with roots on $|z|_p = 1$. (We discuss another modification in Subsection 2.4.) Namely,

Theorem 2.5. Let $f(t) \in \mathbb{Z}[t^{\pm 1}]$ $(\neq 0)$ which may have roots on $|z|_p = 1$. If we modify the definition of the p-adic Mahler measure (Definition 2.3) as

$$\log \mathrm{M}_p(f(t)) = \lim_{n \in \mathbb{N}} \frac{1}{n} \sum_{\zeta^n = 1; f(\zeta) \neq 0} \log |f(\zeta)|_p,$$

then Jensen's formula (Theorem 2.3) still holds.

We give the proof in the next subsection. It is nontrivial if $f(t)$ has a root on $|z|_p = 1$ which is not a root of unity. (We have $|1 + p|_p = 1$ for instance.) Since there is no analogous notions of $\arg \alpha$ and the geometric trigonometric functions for p-adic numbers, we cannot directly follow the proof in [GAS91]. We need to give an alternative framework using properties of roots of unity and the strong triangle inequality.

By the theorem above, we obtain a p-adic analogue of the asymptotic formula:

Proposition 2.6. If $0 \neq f(t) \in \mathbb{Z}[t]$, then

$$\lim_{n \in \mathbb{N}; \text{value} \neq 0} |R(f(t), \nu_n(t))|_p^{1/n} = \mathrm{M}_p(f(t)).$$

We note that the p-adic Mahler measure of a polynomial can be calculated easily from its coefficients (e.g. the argument in the end of [LW88]):

Proposition 2.7. If $f(t) = \sum_i a_i t^{d-i} \in \mathbb{C}_p[t^{\pm 1}]$, then $\mathrm{M}_p(f(t)) = \max\{|a_i|_p\}_i$ holds. In other words, the p-adic Mahler measure and the Gauss norm of $f(t)$ coincide.

Proof. Let the roots of $f(t)$ be indexed by natural numbers as α_i in a descending order with respect to the p-adic norm. If r is the largest i with $|\alpha_i|_p > 1$, then we have $|a_r/a_0|_p = |\alpha_1 \cdots \alpha_r + \text{smaller terms}|_p = |\alpha_1 \cdots \alpha_r|_p$. □

2.3. **Proofs.** To begin with, we recall properties of the p-adic norm $|\cdot|_p$. It defines a ultra-metric, that is, it satisfies the strong triangle inequality $|x-y|_p \leq \max\{|x|_p, |y|_p\}$ for every $x, y \in \mathbb{C}_p$. As a general property of ultra-metric spaces, we have the Krull sharpening: If $|x|_p \neq |y|_p$, then $|x-y|_p = \max\{|x|_p, |y|_p\}$ holds.

Next, we recall several properties of numbers on the p-adic unit circle.

Lemma 2.8. If $\alpha \in \mathbb{C}_p$ satisfies $|\alpha|_p = 1$ and $|\alpha - 1|_p < 1$, and if m is prime to p, then $|\alpha^m - 1|_p = |\alpha - 1|_p$ holds.

Proof. Indeed, we have $|\alpha^m - 1|_p = |\alpha - 1|_p |\alpha^{m-1} + \cdots + 1|_p = |\alpha - 1|_p |(\alpha^{m-1} - 1) + \cdots + (\alpha - 1) + m|_p = |\alpha - 1|_p |(\alpha - 1)(\alpha^{m-2} + \cdots + 1) + m|_p$. Since $|\alpha - 1|_p < 1$, $|\alpha^{m-2} + \cdots + 1|_p \leq \max\{|\alpha^i|_p\}_i = 1$, and $|m|_p = 1$, we have $|(\alpha - 1)(\alpha^{m-2} + \cdots + 1) + m|_p = 1$. □

Lemma 2.9. Let ζ be a primitive n-th root of unity and suppose $n = mp^r$ with $p \not| m$. If $n = 1$, then $|1 - \zeta|_p = 0$. If $m = 1$ and $r > 0$, that is, if ζ is a primitive p-power-th root of unity with $\zeta \neq 1$, then $|1 - \zeta|_p = p^{-1/(p-1)p^{r-1}}$ holds. If otherwise, then $|1 - \zeta|_p = 1$.

Proof. We have $\prod_{\zeta^n = 1, \zeta \neq 1} |1 - \zeta|_p = |R(\nu_n(t), t - 1)|_p = |\nu_n(1)|_p = |n|_p = |p^r|_p = p^{-r}$ and $|1 - \zeta|_p \leq 1$. Since the set of primitive p^r-th roots of unity is closed by non-p powers, the assertion follows from the previous lemma. □

Lemma 2.10. If $\alpha \in \mathbb{C}_p$ satisfies $|\alpha|_p = 1$, then there exist a unique N with $p \not| N$ and a unique primitive N-th root ξ of unity satisfying $|\alpha - \xi|_p < 1$.

Proof. We consider the valuation ring $\mathcal{O}_p = \{z \in \mathbb{C}_p \mid |z|_p \leq 1\}$ and its maximal ideal $\mathfrak{m}_p = \cup_{n \in \mathbb{N}_{>0}} (p^{1/n})$. Since $|\alpha|_p = 1$, we have $\alpha \in \mathcal{O}_p$. We consider $\bmod \mathfrak{m}_p : \mathcal{O}_p \twoheadrightarrow \overline{\mathbb{F}}_p$. Elements of $\overline{\mathbb{F}}_p^\times$ are non-p-power-th roots of unity, and the restriction map $\bmod \mathfrak{m}_p : \{z \in \mathbb{C}_p \mid \text{non-}p\text{-power-th roots of unity}\} \subset \mathcal{O}_p \to \overline{\mathbb{F}}_p^\times$ is a bijection.

Thus there exists a unique non-p-power-th root of unity $\xi \in \mathbb{C}_p$ satisfying $\xi \bmod \mathfrak{m}_p = \alpha \bmod \mathfrak{m}_p$, namely, $|\alpha - \xi|_p < 1$. (This ξ is the image of $\alpha \bmod \mathfrak{m}_p$ under the unique Teichmüller lift $\omega : \overline{\mathbb{F}}_p \hookrightarrow \mathbb{C}_p$.) □

Lemma 2.11. If $\alpha \in \mathbb{C}_p$ satisfies $|\alpha|_p = 1$ and $\alpha \neq 1$, then $\lim_{n \in \mathbb{N}; \alpha^n \neq 1} |\alpha^n - 1|_p^{1/n} = 1$ holds.

Proof. By the previous lemma, we have a unique non-p-power-th root ξ of unity with $|\alpha - \xi|_p < 1$. Suppose that ξ is a primitive N-th root of unity. Then $N \neq 1$. Suppose $\alpha^n \neq 1$. Note that $|\alpha^n - 1|_p = \prod_{\zeta^n = 1} |\alpha - \zeta|_p = \prod_{\zeta^n = 1} |(\alpha - \xi) + (\xi - \zeta)|_p$ holds.

If $N \not| n$ and $\zeta^n = 1$, then $|\xi - \zeta|_p = |1 - \zeta/\xi|_p = 1$. Hence for any $n \in \mathbb{N}$ with $\alpha^n \neq 1$, we have $|\alpha^n - 1| = 1$. If $N | n$, then we have $\{|\xi - \zeta|_p \mid \zeta^n = 1\} = \{|1 - \zeta|_p \mid \zeta^n = 1\}$. The product of all the non-zero elements of this set is $|n|_p$. If ζ runs through primitive p^r-th roots of unity for $r \in \mathbb{N}$, then $|1 - \zeta|_p$ increase with respect to r and converges to 1 as $r \to \infty$. If ζ is another root of unity with $\zeta \neq 1$, then $|1 - \zeta|_p = 1$. Hence we have $|\alpha - \zeta|_p = |\xi - \zeta|_p$ for almost all the roots ζ of unity.

Thus we have $|\alpha^n - 1|_p = \prod_{\zeta^n = 1} |\alpha - \zeta|_p = \bullet \times |n|_p$, where $\bullet$ lies in a finite set with $\bullet \neq 0$. Since $\lim_{n \in \mathbb{N}} |n|_p^{1/n} = 1$, we obtain $\lim_{n \in \mathbb{N}; \alpha^n \neq 1} |\alpha^n - 1|_p^{1/n} = 1$. □

Now let $f(t) = a_0 t^{d-l} \prod_i (t - \alpha_i) \in \mathbb{C}_p[t]$. In order to prove Theorem 2.5, it is sufficient to prove

$$\lim_{n\in\mathbb{N}} \prod_{\zeta^n=1; f(\zeta)\neq 0} |f(\zeta)|_p^{1/n} = |a_0|_p \prod_i \max\{|\alpha_i|_p, 1\}$$

where ζ runs through roots of unity in $\mathbb{C}_p$. Since the both side of this equality is multiplicative with respect to f, it is sufficient to prove the assertion on each factor $t - \alpha_i$. We prove the following lemma:

Lemma 2.12. For any element $\alpha \in \mathbb{C}_p$, the equality $\lim_{n\in\mathbb{N}} \prod_{\zeta^n=1; f(\zeta)\neq 0} |\zeta - \alpha|_p^{1/n} = \max\{|\alpha|_p, 1\}$ holds.

Proof. Suppose that α is not a root of unity. Then we have

$$\prod_{\zeta^n=1; f(\zeta)\neq 0} |\zeta - \alpha|_p = \frac{\prod_{\zeta^n=1} |\zeta - \alpha|_p}{\prod_{\zeta^n=1; f(\zeta)=0} |\zeta - \alpha|_p}.$$

The denominator lies in a finite set independent of n. Hence $(\text{denominator})^{1/n} \to 1$ as $n \to \infty$. If $|\alpha|_p \neq 1$, then $(\text{numerator})^{1/n} = |\alpha^n - 1|_p^{1/n} = \max\{|\alpha|_p^n, 1\}^{1/n} = \max\{|\alpha|_p, 1\}$ holds. If $|\alpha|_p = 1$, then the previous lemma assures $\lim_n |\alpha^n - 1|_p^{1/n} = 1$.

Suppose instead that α is a root of unity. Then we have

$$\prod_{\zeta^n=1; f(\zeta)\neq 0} |\zeta - \alpha|_p = \frac{\prod_{\zeta^n=1; \zeta\neq\alpha} |\zeta - \alpha|_p}{\prod_{\zeta^n=1; f(\zeta)=0, \zeta\neq\alpha} |\zeta - \alpha|_p}.$$

We have $(\text{denominator})^{1/n} \to 1$ as $n \to \infty$. For n's with $\alpha^n \neq 1$, we have $(\text{numerator}) = |\alpha^n - 1|_p$. The previous lemma assures $(\text{numerator})^{1/n} \to 1$ as $n \to \infty$. For n's with $\alpha^n = 1$, we have $(\text{numerator}) = \prod_{\zeta^n=1, \zeta\neq 1} |\zeta - 1|_p = |n|_p$. Since $1/n \leq |n|_p \leq 1$, we have $(\text{numerator})^{1/n} \to 1$ as $n \to \infty$. □

Now we have proved Theorem 2.5.

2.4. **Alternative modification of the integral path.** The Euclidean Mahler measure of $f(t) \in \mathbb{C}[t^{\pm 1}]$ satisfying Jensen's formula was defined as the limit of the integral along a path $\gamma : S^1 \to \mathbb{C}$ as γ uniformly approaches the unit circle. [EW99]

Here we give an analogous argument for our p-adic Mahler measure of $f(t) \in \mathbb{C}_p[t^{\pm 1}]$. We consider the image of roots of unity under a function $g : \mathbb{C}_p \to \mathbb{C}_p$ and let g approach id uniformly with respect to the p-adic topology. Namely, for each function g with $f(g(\zeta)) \neq 0$ for any root ζ of unity, we put $\log \mathrm{M}_{p,g}(f(t)) := \lim_{n\in\mathbb{N}} \frac{1}{n} \sum_{\zeta^n=1} \log |f(g(\zeta))|_p$, where the limit is taken with respect to the Euclidean topology. For each $\varepsilon > 0$, we consider g's with $\sup\{|g(\zeta) - \zeta|_p \mid \zeta \in (\text{roots of unity})\} < \varepsilon$ and let $\varepsilon \to 0$. (Such g exists for each ε, e.g., $g(z) = z + \varepsilon/2$.) Then we have

Proposition 2.13. For any function g sufficiently close to id, the limit $\mathrm{M}_{p,g}(f(t))$ exists and the value is independent of g. The limit $\lim_{g\to\mathrm{id}} \mathrm{M}_p(f(g(t)))$ exists and coincides with $\mathrm{M}_p(f(t))$ satisfying Jensen's formula.

Proof. Since $\log|\bullet|_p$ satisfies the logarithm laws, it is sufficient to consider $f(t) = t - \alpha$ for $\alpha \in \mathbb{C}_p$. If $|\alpha|_p \neq 1$, then trivial. If $|\alpha|_p = 1$, then we have a unique non-p-power-th root ξ of unity satisfying $|\alpha - \xi|_p < 1$. Assume $\sup\{|g(\zeta) - \zeta|_p \mid \zeta \in (\text{roots of unity})\} < \varepsilon < 1$. If $\zeta^n = 1$, then we have $|g(\zeta) - \alpha|_p = |(g(\zeta) - \zeta) + (\zeta - \xi) + (\xi - \alpha)|_p$. By a similar argument to Lemma 2.11, we obtain $\lim_{n\in\mathbb{N}} \prod_{\zeta^n=1} |g(\zeta) - \alpha|_p = 1$. Therefore we have $\mathrm{M}_{p,g}(f(t)) = \max\{|\alpha|_p, 1\}$. □

3. p-adic Mahler measure and $\mathbb{Z}$-covers of links

3.1. **$\mathbb{Z}$-covers and M_p.** Let M be a $\mathbb{Q}\mathrm{HS}^3$ and let L be a d-component link in M. Let $h_\infty : X_\infty \to X = M - L$ be a $\mathbb{Z}$-cover, which we call a *$\mathbb{Z}$-cover over* (M, L), with a generator t of the deck transformation group $t^{\mathbb{Z}}$. Then the Alexander polynomial $A_L(t) \in \mathbb{Z}[t] \subset \mathbb{Z}[t^{\mathbb{Z}}]$ is a generator of the principal ideal of the Fitting ideal $\mathrm{Fitt}_{\mathbb{Z}[t^{\mathbb{Z}}]} H_1(X_\infty)$ of the Alexander module over $\mathbb{Z}[t^{\mathbb{Z}}]$, as explained in Section 1.

Example 3.1. Suppose that every component L_i of L is null-homologous, and let $t_i \in H_1(X)$ denote the meridian of each L_i. Then a standard $\mathbb{Z}$-cover called *the total linking number cover* (*TLN-cover*) *over* (M, L) is defined by the surjective homomorphism $\tau : H_1(X, \mathbb{Z}) \twoheadrightarrow \mathbb{Z}$; $\forall t_i \mapsto 1$. If $A_L(t) \neq 0$, then we have $\mathrm{Fitt}_{\mathbb{Z}[t^{\mathbb{Z}}]} H_1(X_\infty) = (A_L(t))$ by [KM08, Lemma 3.1].

If $M = S^3$, then the multivariable Alexander polynomial $\Delta_L(t_1, \dots, t_d)$, the reduced Alexander polynomial $\Delta_L(t) = \Delta_L(t, \dots, t)$, and the Hosokawa polynomial $H_L(t)$ of L are defined. For the TLN-cover over (S^3, L), we have

$$A_L(t) = (t-1)^{d-1} H_L(t) = \begin{cases} \Delta_L(t) & \text{if } d = 1, \\ (t-1)\Delta_L(t) & \text{if } d > 1. \end{cases}$$

For each $n \in \mathbb{N}_{>0}$, let $M_n \to M$ denote the branched $\mathbb{Z}/n\mathbb{Z}$-cover obtained as the Fox completion of the $\mathbb{Z}/n\mathbb{Z}$-subcover of h_∞. Then the orders of groups and the cyclic resultants of $A_L(t)$ satisfy the following well-known formula:

Proposition 3.2. Let the notation be as above. Then there exists some finite set $\mathcal{C}$ of $\mathbb{N}$ such that for any $n \in \mathbb{N}_{>0}$

$$|H_1(M_n)| = |R(A_L(t), \nu_n(t))| \times C$$

holds for some $C \in \mathcal{C}$.

This proposition follows from Sakuma's theorem ([Sak81, Theorem 3]) or Porti's theorem ([Por04, Theorem 1.1]) by a similar argument to [KM08, Proof of Theorem 2.1]. It was initially proved for a knot L in $M = S^3$ by Fox ([Fox56], [Web79]) and for a link L in $M = S^3$ by Mayberry–Murasugi [MM82].

By Propositions 2.1, 2.6, and Theorem 3.2, we obtain the following asymptotic formulae:

Theorem 3.3. Let the notation be as above. If $A_L(t) \neq 0$, then

$$\lim_{n\in\mathbb{N};\text{value}\neq 0} |H_1(M_n)|^{1/n} = \mathrm{M}(A_L(t)),$$
$$\lim_{n\in\mathbb{N};\text{value}\neq 0} ||H_1(M_n)||_p^{1/n} = \mathrm{M}_p(A_L(t)).$$

The Euclidean formula for knots in S^3 was proved by [GAS91], [Ril90]. Riley also gave an estimation of $||H_1(M_n)||_p^{-1}$ from above.

We denote $p = \infty$ for Euclidean objects. For $f(t) \in \mathbb{Z}[t^{\pm 1}]$, the set of p-adic Mahler measures $\mathrm{M}_p(f(t))$ for $p < \infty$ do not tell that for $p = \infty$. The set of those for $p \leq \infty$ would have some meaning.

Example 3.4. We use the notation in the Rolfsen table ([Rol76]).

(1) If $K = 4_1$ (figure 8-knot), then the Alexander polynomial $\Delta_K(t) = t^2 - 3t + 1$ satisfies $\mathrm{M}(\Delta_K(t)) = \frac{3+\sqrt{5}}{2}$, $\mathrm{M}_p(\Delta_K(t)) = 0$ $(p < \infty)$.

(2) If $L = 4_1^2$ (Solomon's knot, sigillum Salomonis), then one of the reduced Alexander polynomial $\Delta_L(t, t^{-1}) = 2$ satisfies $\mathrm{M}(2) = 0, \mathrm{M}_2(2) = 1/2, \mathrm{M}_p(2) = 0$ $(p \neq 2)$. The same values are obtained for $A_L(t) = 2(t-1)$.

Remark 3.5. ($\mathbb{Z}^d$-covers, non-$\mathbb{Q}\mathrm{HS}^3$-cases) We have an asymptotic formula for $\mathbb{Z}^d$-covers with use of the multivariable Alexander polynomials and the Mahler measures. In addition, even if M_n's are not necessarily $\mathbb{Q}\mathrm{HS}^3$, we have an asymptotic formula of the torsion subgroups $H_1(M_n)_{\mathrm{tor}}$ with use of the higher Alexander polynomials and the Mahler measures ([SW02], [Le14]). Note that sometimes we obtain special values of zeta functions, L-functions, and multiple zeta functions as the Mahler measures of multivariable polynomials ([Smy81], [Lal03]). An asymptotic formula with use of the hyperbolic volumes is known as the Bergeron–Venkatesh conjecture and Le's theorem ([BV13], [Le14]). We expect p-adic analogues of them.

3.2. **Iwasawa μ_p-invariant and M_p.** Let M be a $\mathbb{Q}\mathrm{HS}^3$, let L be a link in M, and let $\{h_{p^r} : M_{p^r} \to M\}_r$ be an inverse system of branched $\mathbb{Z}/p^r\mathbb{Z}$-covers over (M, L). (For such an object, see Subsection 5.1 also.) Let t denote the topological generator of the inverse limit of the deck transformation groups $\mathbb{Z}/p^r\mathbb{Z}$ corresponding to 1. Then the Iwasawa module $\mathcal{H}_p := \varprojlim H_1(M_{p^r}, \mathbb{Z}_p)$ is a finitely generated module over $\mathbb{Z}_p[[t^{\mathbb{Z}_p}]] \cong \mathbb{Z}_p[[T]]$. By [Uek16a, Theorem 4.17], M_{p^r}'s are $\mathbb{Q}\mathrm{HS}^3$'s if and only if its characteristic polynomial in $\mathbb{Z}_p[[t^{\mathbb{Z}_p}]](\cong \mathbb{Z}_p[[T]])$ does not vanish at any p-power-th root of unity.

The following formula is an analogue of Iwasawa's class number formula ([Iwa59]), which was initially given by [HMM06] and generalized by [KM08] and [Uek16a]:

Proposition 3.6 (Iwasawa type formula)**.** Let the notation be as above. If M_{p^n}'s are $\mathbb{Q}\mathrm{HS}^3$'s, then there exist some $\lambda_p, \mu_p \in \mathbb{N}, \nu_p \in \mathbb{Z}$ called the Iwasawa invariants such that for any $r \gg 0$ in $\mathbb{N}$

$$||H_1(M_{p^r})||_p^{-1} = p^{\lambda_p r + \mu_p p^r + \nu_p}$$

holds.

Suppose that the $\mathbb{Z}/p^r\mathbb{Z}$-covers are obtained from a $\mathbb{Z}$-cover over (M, L) with the Alexander polynomial $A_L(t)$. Comparing with the results in the previous section, we obtain

Proposition 3.7.

$$\log \mathrm{M}_p(A_L(t)) = -\mu_p \log p.$$

In this sense, the Iwasawa type formula is a p-adic refinement of the asymptotic formula (Theorem 3.3) with use of M_p.

Remark 3.8. We review what we have done for $\mathbb{Z}$-covers of S^3 branched over links L. By the Mayberry–Murasugi formula and the definition of M_p imitating the Shnirel'man integral, if $A_L(t)$ does not vanish on the p-adic unit circle, then we have $\lim_{n \in \mathbb{N}; \gcd(p,n)=1} ||H_1(M_n)||_p = \mathrm{M}_p(A_L(t))$. By the Iwasawa type

formula, if $A_L(t)$ does not vanish at any p-power-th root of unity, then we have $\lim_{n=p^r} ||H_1(M_n)||_p = p^{-\mu_p}$. Note that the intersection of their domains of n is empty. By Jensen's formula, the formula $\mathrm{M}_p(A_L(t)) = \max\{|\text{coefficients}|_p\}$, and the p-adic Weierstrass preparation theorem, if $A_L(t)$ has no root on $|z|_p = 1$, then $\log \mathrm{M}_p = -\mu_p \log p$ immediately follows. We removed the assumption on $A_L(t)$ by extending the definition of M_p.

We have $\mathrm{M}_p(A_L(t)) = \max\{|\text{coefficients}|_p\}$. If $L = K$ is a knot, then $A_L(t) = \Delta_K(t)$ satisfies $\Delta_K(1) = \pm 1$ and its is primitive. Thus for each $n < \infty$, we have $\mathrm{M}_p(A_L(t)) = 1$ and $\mu_p = 0$. In regard to $d \geq 2$-component link L, various examples with $\mathrm{M}_p(A_L(t)) < 1$ and $\mu_p > 0$ are given ([KM08], [KM13], [Uek16b]).

Example 3.9. We list up all the examples of TLN-covers with non-trivial μ_p obtained from 2-component links L in the Rolfsen table ([Rol76]). We have $A_L(t) = (t-1)\Delta_L(t^{\epsilon_1}, t^{\epsilon_2}), \epsilon_i \in \{\pm 1\}$. We denote $f \doteq g$ if f and g coincide up to multiplication by units of $\mathbb{Z}[t^{\pm 1}]$.

(1) $L = 4_1^2$. $\Delta_L(x, y) = 1 + xy$, $\Delta_L(t, t^{-1}) = 2$, $\mu_2 = 1$.
(2) $L = 6_1^2$. $\Delta_L(x, y) = 1 + xy + x^2y^2$, $\Delta_L(t, t^{-1}) = 3$, $\mu_3 = 1$.
(3) $L = 6_3^2$. $\Delta_L(x, y) = 2 - x - y + 2xy$, $\Delta_L(t, t) = 2(t^2 - t + 1)$, $\mu_2 = 1$.
(4) $L = 7_3^2$. $\Delta_L(x, y) = 1 - x - y + xy$, $\Delta_L(t, t) \doteq \Delta_L(t, t^{-1}) \doteq 4(t^2 - 1)^2$, $\mu_2 = 2$.

3.3. **p-adic entropy h_p and M_p.** Noguchi studied $\mathbb{Z}$-covers of S^3 branched over knots ([Nog07]). Let t denote the meridian action on the ($\mathbb{Q}$-)Alexander module $H_1(X_\infty, \mathbb{Q}) \cong \mathbb{Q}^l$, and let $\widehat{t}$ denote the solenoidal system obtained as the Pontrjagin dual of t. He applied the result of Lind and Ward ([LW88]) to $\widehat{t}$ and calculated the topological entropy h_{top} of $\widehat{t}$, which coincides with Bowen's measure theoretic entropy.

Giordano-Bruno and Virili ([GBV15]) studied the algebraic entropies h_{alg} of dynamical systems of $\mathbb{Q}$-linear spaces. They gave a formula parallel to that of [LW88], so that h_{alg} of t coincides with h_{top} of $\widehat{t}$ above. In what follows, we just call them the entropy.

These entropies have their origin in the Ergodic theory. They were introduced by Adler, Kronheim, and McAndrew ([AKM65]). Various generalization and duality have been studied ([DGB14] for instance).

Results of [LW88] and [GBV15] called the Yuzvinski formula or the Kolmogorov–Sinai formula can be stated with use of our p-adic Mahler measure as follows:

Proposition 3.10. Let $\varphi \in \mathrm{GL}(l, \mathbb{Q})$ with $l \in \mathbb{N}$ and let $B(t) \in \mathbb{Q}[t]$ denote its (monic) characteristic polynomial. Then the algebraic entropy of $\varphi \curvearrowright \mathbb{Q}^l$ and the topological entropy of its Pontrjagin dual are given by

$$h = \sum_{p \leq \infty} h_p, \quad h_p = \log \mathrm{M}_p(B(t)),$$

where h_p denotes the p-adic entropy of $\varphi \curvearrowright \mathbb{Q}^l \otimes \mathbb{Q}_p \cong \mathbb{Q}_p^l$ or its Pontrjagin dual for $p \leq \infty$.

In addition, suppose $B(t) = \sum_i b_i t^{d-i} \in \mathbb{Q}[t]$. Then $\sum_{p<\infty} h_p = \log s$ hols for $s = \mathrm{lcm}\{b_i\}_i$, and the primitive polynomial $sB(t) \in \mathbb{Z}[t]$ satisfies $h = \log \mathrm{M}(sB(t))$.

Now let L be a link in S^3 and let $h_\infty : X_\infty \to X = S^3 - L$ be a TLN-$\mathbb{Z}$-cover with the Alexander polynomial $A_L(t) = \sum_{0 \leq i \leq l} a_i t^{d-i} \in \mathbb{Z}[t]$ with $l, d \in \mathbb{N}$. Then

we have $H_1(X_\infty;\mathbb{Q}) \cong \mathbb{Q}^l$ as linear space over $\mathbb{Q}$ and the characteristic polynomial of the meridian action on $H_1(X_\infty;\mathbb{Q})$ is given by the associated monic polynomial $A_L(t)/a_0$. Hence we have

Proposition 3.11. The entropy of the t action on $H_1(X_\infty;\mathbb{Q})$ is given by

$$h = \sum_{p\le\infty} h_p, \quad h_p = \log \mathrm{M}_p(A_L(t)/a_0).$$

By the product formula $\prod_{p\le\infty}|a_0|_p = 1$, we have $h = \sum_{p\le\infty}\log \mathrm{M}_p(A_L(t))$ also. In addition, the primitive polynomial $A_L(t)/\gcd\{a_i\}_i \in \mathbb{Z}[t]$ satisfies $h = \mathrm{M}(A_L(t)/\gcd\{a_i\}_i)$.

If L is a knot, then $A_L(t) = \Delta_L(t)$ is primitive, and similar results hold for the Alexander module $H_1(X_\infty)$ over $\mathbb{Z}$ ([Nog07, Theorem 1]). If L is a general link, then $A_L(t)$ is not necessarily primitive. The entropy $h_\mathbb{Z}$ of the meridian action on $H_1(X_\infty)$ over $\mathbb{Z}$ is given by $h_\mathbb{Z} = \log \mathrm{M}(A_L(t))$ and satisfies $h_\mathbb{Z} = h + \log\gcd\{a_i\}_i$.

Example 3.12. In the Rolfsen table ([Rol76]), the first example with nontrivial entropy h is $K = 4_1$ (figure 8-knot). We have $\Delta_K(t) = t^2 - 3t + 1$, $h = \log \mathrm{M}_\infty(\Delta_K(t)) = \log\frac{3+\sqrt{5}}{2}$. We list up all the 2-component links L with nontrivial h in the table:

(1) $L = 6_2^2$. $\Delta_L(x,y) = x+y-xy+x^2y+xy^2$, $\Delta_L(t,t) \doteq \Delta_L(t,t^{-1}) \doteq 2t^2-t+2$, $h = h_2 = \log 2$.

(2) $L = 6_3^2$. $\Delta_L(x,y) = 2 - x - y + 2xy$, $\Delta_L(t,t^{-1}) \doteq t^2 - 4t + 1$, $h = h_\infty = \log(2+\sqrt{3})$.

(3) $L = 7_1^2$. $\Delta_L(x,y) = 1 - x - y + xy - x^2y - xy^2 + x^2y^2$. (i) $\Delta_L(t,t) = t^4 - 2t^3 + t^2 - 2t + 1$, $h = h_\infty = \log\frac{1+\sqrt{2}+\sqrt{2\sqrt{2}-1}}{2}$. (ii) $\Delta_L(t,t^{-1}) \doteq 2t^2 - 3t + 2$, $h = h_2 = \log 2$.

(4) $L = 7_2^2$. $\Delta_L(x,y) = 1-x-y+3xy-x^2y-xy^2+x^2y^2$, $\Delta_L(t,t^{-1}) \doteq 2t^2-5t+2 = 2(t-2)(t-\frac{1}{2})$, $h_\infty = 2$, $h_2 = \log|1/2|_2 = \log 2$, $h = h_\infty + h_2 = \log \mathrm{M}(2t^2-5t+2) = 2\log 2$.

3.4. **h_p, μ_p, and the leading coefficient a_0.** We continue to study a TLN-$\mathbb{Z}$-cover of (S^3, L). The leading coefficient a_0 of $A_L(t)$ has geometric information. For instance, if L is fibered, then $a_0 = 1$ holds (e.g., [Hil12, Theorem 5.12]).

Now let p be a prime number again. Then $h_p = \log \mathrm{M}_p(A_L(t)/a_0)$ measures the ratio between the monic and primitive polynomials associated to $A_L(t)$, while $\mathrm{M}_p(A_L(t)) = p^{-\mu_p}$ measures the ratio between $A_L(t)$ itself and the primitive one.

Since $\mathrm{M}_p(A_L(t)/a_0) = \mathrm{M}_p(A_L(t))/|a_0|_p$, we obtain a balance formula among h_p, μ_p, and a_0:

Proposition 3.13.

$$-\log|a_0|_p = h_p + \mu_p \log p.$$

Example 3.14. We have only one example of 2-component link whose h_p, μ_p are both nontrivial in the Rolfsen table ([Rol76]). It is $L = 9_{23}^2$. We have $\Delta_L(x,y) = x^2 - 2xy + y^2x^3 - 4x^2y - 4xy - y^3 + x^3y + 2x^2y^2 + xy^3$, $\Delta_L(t,t) = 4t^2 - 10t + 4 = 2(2t^2 - 5t + 2) = 2(2-t)(1-2t)$. $|a_0|_2 = 1/4$. $h_\infty = \log 2$, $h_2 = \log|1/2|_2 = \log 2$, $h = h_\infty + h_2 = \log \mathrm{M}_\infty(2t^2 - 5t + 2) = 2\log 2$. $\mathrm{M}_2(\Delta_L(t,t)) = |2|_2 = 1/2$, $\mu_2 = 1$.

If $A_L(t) = a_0 \prod_i (t - \alpha_i)$, then by Theorems 2.3, 2.5, and Proposition 3.11, we have $h_p = \sum_{|\alpha_i|_p > 1} \log |\alpha_i|_p$. Note the product formula $\prod_{p<\infty} |a_0|_p = 1/|a_0|$. Then the previous proposition gives

Corollary 3.15.

$$\log |a_0| = \sum_{p<\infty} \sum_{|\alpha_i|_p > 1} \log |\alpha_i|_p + \sum_{p<\infty} \mu_p \log p.$$

This formula generalizes [Nog07, Corollary 4]. Indeed, if L is a knot, then $\mu_p = 0$ for any $p < \infty$, and hence $\log |a_0| = \sum_{p<\infty} \sum_{|\alpha_i|_p > 1} \log |\alpha_i|_p$ $(= \sum_{p<\infty} h_p)$ holds.

4. Purely p-adic notions

4.1. **Besser–Deninger m_p.** The notion of the p-adic logarithm $\log_p : \mathbb{C}_p^* \to \mathbb{C}_p$ is known ([Was97, Chapter 5]). It is first defined on the unit disc $|z-1|_p < 1$ by the power series $\log_p(1+z) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1} z^n}{n}$, and then uniquely extended to whole $\mathbb{C}_p^*$ by the p-adic analytic continuation under the normalization $\log_p p = 0$. It is sometimes called the Iwasawa p-adic logarithm. Besser and Deninger introduced the *purely* p-adic logarithmic Mahler measure m_p for p-adic analytic functions ([BD99]). It is different from ours, and is defined by the Shnirel'man integral ([Shn38]):

Definition 4.1. For a p-adic analytic functions $f(z)$ with no zero on $|z|_p = 1$, we put

$$m_p(f(z)) := \int_{|z|_p=1} \log_p f(z) \frac{dz}{z} := \lim_{n \in \mathbb{N}; \gcd(n,p)=1} \sum_{\zeta^n = 1} \log_p f(z),$$

where the limit is taken with respect to the p-adic topology.

It satisfies Jensen's formula for Laurent polynomials:

Proposition 4.2 (Jense's formula)**.** If $f(t) \in \mathbb{C}_p[t^{\pm 1}]$ with no root on $|z|_p = 1$, then the limit in Definition 4.1 exists. If $f(t) = \sum_{0 \le i \le l} a_i t^{d-i} = a_0 t^{d-l} \prod_{0 \le i \le l} (t - \alpha_i)$ $(a_i, \alpha_i \in \mathbb{C}_p, a_0 a_l \neq 0)$, then we have

$$\begin{aligned} m_p(f(t)) &= \log_p a_l - \sum_{0 < |\alpha_i|_p < 1} \log_p \alpha_i \\ &= \log_p a_0 + \sum_{|\alpha_i|_p > 1} \log_p \alpha_i. \end{aligned}$$

Since $\log_p(z)$ does not vanish on $|z|_p = 1$, we have no natural generalization of m_p for $f(z)$ with zeros on $|z|_p = 1$.

Remark 4.3. In the Shnirel'man integral, we have no term corresponding to $2\pi\sqrt{-1}$ in the complex case.In [Mih12], its *denormalization* and generalization are given, in which the corresponding terms called periods in $\mathbb{B}_{\mathrm{dR}}$ appear.

By the definition of the Shnirel'man integral, an analogue of the asymptotic formula of resultants (Proposition 2.1, 2.6) immediately follows:

Proposition 4.4. If $f(z) \in \mathbb{C}_p[t]$ with no root on $|z|_p = 1$, then

$$\lim_{n \in \mathbb{N}; (n,p)=1} \frac{1}{n} \log_p |R(f(t), t^n - 1)| = m_p(f(t)).$$

Since $R(f(t), t^n - 1) = R(f(t), \nu_n(t)) \times f(1)$, if $f(1) \neq 0$, then the convergence of $\{\frac{1}{n} \log_p |R(f(t), \nu_n(t))|\}_n$ is equivalent to that of $\{\frac{1}{n} \log_p |f(1)|\}_n$, and hence to $\log_p |f(1)| = 0$.

We apply their theory to the TLN-$\mathbb{Z}$-cover of S^3 branched over a d-component link L. Put $\Lambda = \mathbb{Z}[t^{\pm 1}]$. We use the notation in the previous section. We have $A_L(t) = (t-1)^d H_L(t)$. The Alexander module admits a natural direct sum decomposition $H_1(X_\infty) \cong \mathbb{Z}^{d-1} \oplus H_1'(X_\infty)$ of $\mathbb{Z}[t^{\mathbb{Z}}]$-modules with $\mathrm{Fitt}_{\mathbb{Z}[t^{\mathbb{Z}}]} H_1'(X_\infty) = (H_L(t))$.

If $d = 1$, then $A_L(t)$, $H_L(t)$ and the Alexander polynomial $\Delta_L(t)$ coincide. Since $A_L(1) = \pm 1$, by Proposition 3.2, we have $|H_1(M_n)| = |R(A_L(t), \nu_n(t)))| = |R(A_L(t), t^n - 1)|$. We obtain an analogue of Theorem 3.3 with respect to the p-adic norm: If $A_L(t)$ has no root on $|z|_p = 1$, then

$$\lim_{n \in \mathbb{N}; (n,p)=1} \frac{1}{n} \log_p |H_1(M_n)| = m_p(A_L(t)).$$

For the cases with $d > 1$, note that the trivial action of t on $\mathbb{Z}$ is not expansive and $\hbar_p$ is not defined. It comes from the fact that $\{(\log_p n)/n\}_n$ is not convergent in $\mathbb{C}_p$. However, for any d, if $H_L(z)$ has no roots on $|z|_p = 1$, then the meridian action t on the Hosokawa module $\Lambda/(H_L(t))$ is expansive, and its purely p-adic entropy is given by $\hbar_p = m_p(H_L(t))$.

For a general $d \in \mathbb{N}_{>0}$, we have $|H_1(M_n)| = |R(A_L(t), \nu_n(t)))| = |R(H_L(t), \nu_n(t))$ $R((t-1)^{d-1}, \nu_n(t))| = |R(H_L(t), t^n - 1)/R(H_L(t), t - 1)| \times n^{d-1} = |R(H_L(t), t^n - 1)/H_L(1)| \times n^{d-1}$. Hence a modified asymptotic formula for links is stated as follows:

Theorem 4.5. Let L be a d-component link in S^3 and suppose that the Hosokawa polynomial $H_L(t)$ has no root on $|z|_p = 1$. Then the TLN-$\mathbb{Z}$-cover over (S^3, L) satisfies

$$\lim_{n \in \mathbb{N}; (n,p)=1} \frac{1}{n} \log_p \frac{|H_1(M_n)||H_L(1)|}{n^{d-1}} = m_p(H_L(t))$$

with respect to the p-adic topology.

4.2. **Deninger's $\hbar_p$.** The topological entropy of a dynamical system (an automorphism φ on a space X) is sometimes calculated by using the numbers of the fixed points (e.g., [EW99, Section 2]):

$$h(\varphi) = \lim_{n \to \infty} \frac{1}{n} \log |\mathrm{Fix}(\varphi^n)|.$$

Deninger defined *purely* p-adic entropy $\hbar_p(\varphi)$ by using the numbers of the fixed points and p-adic logarithm under certain situations:

$$\hbar_p(\varphi) := \lim_{n \to \infty} \frac{1}{n} \log_p |\mathrm{Fix}(\varphi^n)|.$$

A Laurent polynomial f has no zeros on the p-adic unit torus if and only if the associated solenoidal dynamical system is expansive. Under those conditions, we have $\hbar_p = m_p(f)$ ([Den09, Theorem 1]).

We consider the TLN-$\mathbb{Z}$-cover over a d-component link L in S^3 again. We consider the meridian action on the Alexander module $H_1(X_\infty) \cong \mathbb{Z}^{d-1} \oplus \Lambda/(H_L(t))$.

Since $\{(\log_p n)/n\}_n$ is not convergent in $\mathbb{C}_p$, the trivial action of t on $\mathbb{Z}$ is not expansive. Therefore, if $d > 1$, then $\hbar_p$ is not defined for $t \curvearrowright H_1(X_\infty)$. However, we have a formula for a direct summand:

Proposition 4.6. For any d, if $H_L(z)$ has no root on $|z|_p = 1$, then the meridian action on the Hosokawa module $H'_1(X_\infty)$ is expansive, and its purely p-adic entropy is given by $\hbar_p = m_p(H_L(t))$.

Example 4.7. The first example of a knot in Rolfsen's table for which $\hbar_p$ is defined is $L = 5_2$ (the 3-twist knot). We have $A_L(t) = \Delta_L(t) = 2t^2 - 3t + 2$. Its roots α, β satisfy $|\alpha\beta|_2 = 1$ and $|\alpha + \beta|_2 = |3/2|_2 = 2 > 1$. Let $\alpha = (3 - \sqrt{-7})/4$ denote the larger root. Then we have $\hbar_2 = m_2(\Delta_L(t)) = \log_2(3 - \sqrt{-7})$.

Among the examples of links we have seen, $\hbar_p$ is defined only for the following cases with $p = 2$:

Example 3.12 (1) $L = 6_2^2$, $H_L(t) \doteq 2t^2 - t + 2$. Let $(1 - \sqrt{-15})/4$ denote the larger root. Then we have $\hbar_2 = m_2(A_L(t)/2) = \log_2(1 - \sqrt{-15})$.

(3) $L = 7_1^2$. (ii) $H_L(t) \doteq 2t^2 - 3t + 2$. Let $(3 - \sqrt{-7})/4$ denote larger root. Then $\hbar_2 = \log_2(3 - \sqrt{-7})$.

(4) $L = 7_2^2$, $H_L(t) \doteq 2t^2 - 5t + 2 = 2(t - 2)(t - \frac{1}{2})$. $\hbar_2 = 0$.

5. Profinite cyclic covers

We study entropies of the meridian actions on modules associated to profinite cyclic covers.

5.1. **$\mathbb{Z}_p$-covers.** Let L be a link in S^3. A *branched $\mathbb{Z}_p$-cover* over (S^3, L) is an inverse system $\{M_{p^n} \to S^3\}_n$ of branched $\mathbb{Z}/p^n\mathbb{Z}$-covers branched over L. If we fix a $\mathbb{Z}$-cover of $S^3 - L$, then we obtain a branched $\mathbb{Z}_p$-cover from its subcovers. A branched $\mathbb{Z}_p$-cover is not necessarily obtained from a $\mathbb{Z}$-cover, because the image of $\mathbb{Z}^2 \to \mathbb{Z}_p$ is not necessarily sent to $\mathbb{Z} \subset \mathbb{Z}_p$ by automorphisms of $\mathbb{Z}_p$.

We have an isomorphism $\mathbb{Z}_p[[t^{\mathbb{Z}_p}]] \cong \mathbb{Z}_p[[T]]; t \mapsto 1 + T$. The Iwasawa module (pro-p Alexander module) $\mathcal{H}_p := \varprojlim_n H_1(M_{p^n}, \mathbb{Z}_p)$ of a branched $\mathbb{Z}_p$-cover is a finitely generated torsion $\mathbb{Z}_p[[T]]$-module ([Uek16a]). Let $\tau : H_1(S^3 - L) \to \mathbb{Z}_p; t_i \mapsto z_i$ ($z_i \in \mathbb{Z}_p^\times$) be a homomorphism. Then Ker of the composite of τ and the natural surjections $\mathbb{Z}_p \twoheadrightarrow \mathbb{Z}/p^n\mathbb{Z}$ defines an inverse system of $\mathbb{Z}/p^n\mathbb{Z}$-covers, and hence a totally branched $\mathbb{Z}_p$-cover associated to τ. The Fitting ideal of $\mathcal{H}_p$ over $\mathbb{Z}_p[[t^{\mathbb{Z}_p}]]$ is generated by $A_L(t) = (t - 1)\Delta_L(t^{z_1}, \dots, t^{z_d})$.

By the p-adic Weierstrass preparation theorem [Was97, Theorem 7.3], we have $A_L(1 + T) = p^{\mu_p} g(T) u(T)$ for a distinguished polynomial $g(T)$ (that is, a monic polynomial with every lower coefficient divisible by p) and $u(T) \in \mathbb{Z}_p[[T]]^\times$. All the roots of $g(T)$ are on $|z|_p < 1$ and $\lambda_p = \deg(g(T))$ holds. If $u(T)$ has a zero on the domain of convergence, then it is on $|z|_p \geq 1$. We have $\mathcal{H} \otimes \mathbb{Q} \cong \oplus_i \mathbb{Q}_p[[T]]/(g_i(T))$ and $\prod_i g_i(T) = g(T)$ for some $g_i(T)$'s. Thus we have $\mathbb{Q}_p[[T]]/g_i(T) \cong \mathbb{Q}_p^{\deg(g_i)}$ and hence $\mathcal{H}_p \otimes \mathbb{Q} \cong \mathbb{Q}_p^{\lambda_p}$ as linear spaces over $\mathbb{Q}_p$. The characteristic polynomial of the meridian action (the t action) on $\mathcal{H}_p \otimes \mathbb{Q}$ is $g(t - 1) \in \mathbb{Z}[t]$, and hence its entropy is given by

$$h = h_p = \mathrm{M}_p(g(t - 1)) = 0.$$

5.2. **$\widehat{\mathbb{Z}}$-covers.** Next, we consider branched $\widehat{\mathbb{Z}}$-covers. Note that the profinite integer ring admits the decomposition $\widehat{\mathbb{Z}} = \prod_p \mathbb{Z}_p$ by the Chinese remainder theorem. A *branched $\widehat{\mathbb{Z}}$-cover* over (L, S^3) is an inverse system $\{M_n \to S^3\}_n$ of branched $\mathbb{Z}/n\mathbb{Z}$-covers branched over a link. Let $\tau : H_1(S^3 - L) \to \widehat{\mathbb{Z}}; t_i \mapsto z_i,\ z_i \in \widehat{\mathbb{Z}}^\times$ be a homomorphism. Then the kernels of the composites of τ and the natural surjections $\widehat{\mathbb{Z}} \twoheadrightarrow \mathbb{Z}/n\mathbb{Z}$ define an inverse system of branched $\mathbb{Z}/n\mathbb{Z}$-covers, and hence a branched $\widehat{\mathbb{Z}}$-cover associated to τ. We put $A_L(t) := (t-1)\Delta_L(t^{z_1}, \dots, t^{z_d})$ and write $A_L(t) = (t-1)^\delta H_L(t)$, $H_L(t) \in \mathbb{Z}[[t^{\widehat{\mathbb{Z}}}]]$, $H_L(1) \neq 0$. There is a natural isomorphism between $H_1(M_n; \widehat{\mathbb{Z}})$ and the profinite completion $\widehat{H}_1(M_n)$ of $H_1(M_n)$. There is a natural direct sum decomposition $H_1(M_n) \xrightarrow{\cong} \mathbb{Z}/n\mathbb{Z}^{\oplus\delta} \oplus \widehat{H}'_1(M_n)$ with $\delta \in \mathbb{N}$ such that the Fitting ideal of $\widehat{H}'_1(M_n)$ over $\widehat{\mathbb{Z}}[t^{\mathbb{Z}/n\mathbb{Z}}]$ is given by $\mathrm{Fitt}_{\mathbb{Z}[t^{\mathbb{Z}/n\mathbb{Z}}]}\widehat{H}'_1(M_n) = (H_L(t)) \subset \widehat{\mathbb{Z}}[t^{\mathbb{Z}/n\mathbb{Z}}]$.

We define the completed Alexander module by putting $\mathcal{H} := \varprojlim_n H_1(M_n, \widehat{\mathbb{Z}})$. This module is a finitely generated $\widehat{\mathbb{Z}}[[t^{\widehat{\mathbb{Z}}}]]$-module. Indeed, let $z_i = (z_{i,n} \bmod n)_n$ with $z_{i,n} \in \mathbb{Z}$ for each i. Let $h_{\infty,n}$ denote the $\widehat{\mathbb{Z}}$-cover corresponding to the map $\tau_n : H_1(S^3 - L) \mapsto \widehat{\mathbb{Z}}; t_i \mapsto z_{i,n}$ for each n. Then the completed Alexander module $\mathcal{H}_n$ of $h_{\infty,n}$ is a finitely generated $\widehat{\mathbb{Z}}[[t^{\mathbb{Z}}]]$-module by a similar argument to [Uek17, Lemma 11]. Since the ranks of $\mathcal{H}_n$'s are bounded and $\mathcal{H}$ is approximated by $\mathcal{H}_n$'s, $\mathcal{H}$ is a finitely generated $\widehat{\mathbb{Z}}[[t^{\mathbb{Z}}]]$-module. We have a natural direct sum decomposition $\mathcal{H} \cong \widehat{\mathbb{Z}}^{\oplus\delta} \oplus \mathcal{H}'$ such that the Fitting ideal of $\mathcal{H}'$ is $(H_L(t)) \subset \widehat{\mathbb{Z}}[[t^{\mathbb{Z}}]]$.

A branched $\widehat{\mathbb{Z}}$-cover which cannot be obtained from a $\mathbb{Z}$-cover may admit $A_L(t)$ in $\mathbb{Z}[t]$:

Example 5.1. Let $L = K_1 \cup K_2 \cup K_3 \subset S^3$ be a 3-component link with $\Delta_L(x, y, z) = B(xyz) \in \mathbb{Z}[xyz]$. Take $u \in \widehat{\mathbb{Z}}^\times$ with $u \neq \pm 1$. Let t_i denote the meridian of K_i in $H_1(S^3 - L)$ for each i. Consider tha map $\tau : H_1(S^3 - L) \to \widehat{\mathbb{Z}}; t_1 \mapsto 1, t_2 \mapsto u, t_3 \mapsto -u$. Then it defines a branched $\widehat{\mathbb{Z}}$-cover which cannot be obtained from a $\mathbb{Z}$-cover and admits the characteristic polynomial $A_L(t) = (1-t)\Delta_L(t, t^u, t^{-u}) = (1-t)B(t) \in \mathbb{Z}[t]$.

There is only one non-trivial example in the Rolfsen table ([Rol76]): the link $L = 8^3_7$ satisfies $\Delta_L(x, y, z) = 1 - xyz$, $A_L(t) = (1-t)\Delta_L(t^u, t^{-u}, t) = (1-t)^2$.

A $\widehat{\mathbb{Z}}$-cover is approximated by $\mathbb{Z}$-covers, that is, every layer can be obtained as a quotient of a $\mathbb{Z}$-cover. Hence a generalization

$$|\widehat{H}_1(M_n)| = |R(A_L(t), \nu_n(t))|$$

of Fox's formula (Proposition 3.2) holds. Therefore, if $A_L(t) \in \mathbb{Z}[t]$, then for any $p \leq \infty$, generalizations of asymptotic formulae with use of p-adic Mahler measures (Theorems 3.3) hold:

$$\lim_{\text{value}\neq 0} |H_1(M_n)|_p = \mathrm{M}_p(A_L(t)).$$

In addition, if $H_L(t) := A_L(t)/(t-1)^{d-1}$ has no root on roots of unity, then a generalization of the purely p-adic version (Theorem 4.5) for each $p < \infty$ holds:

$$\lim_{n\in\mathbb{N};(n,p)=1} \frac{1}{n} \log_p \frac{|H_1(M_n)|}{n^{d-1}} = m_p(H_L(t)).$$

In the following, we consider a branched $\widehat{\mathbb{Z}}$-cover with $A_L(t) \in \mathbb{Z}[t]$ and study the entropy of the meridian action (t-action). Note the decomposition $\widehat{\mathbb{Z}}[[t^{\widehat{\mathbb{Z}}}]] \cong \prod_p \mathbb{Z}_p[[t^{\widehat{\mathbb{Z}}}]]$. Let $\prod_p \mathbb{Q}_p$ denotes the restricted product of $\mathbb{Q}_p$'s with respect to open subgroups $\mathbb{Z}_p < \mathbb{Q}_p$.

Here we assume $\mathcal{H} \otimes \mathbb{Q} \cong (\prod_p \mathbb{Q}_p)^{\deg(A_L(t))}$. Then for each prime number p, the degree of $A_L(t) \bmod p$ in $\mathbb{F}_p[t]$ coincides with that of $A_L(t) \in \mathbb{Z}[t]$. Indeed, since $\mathcal{H}$ is a quotient of compact Hausdorff group, it has no p-divisible factor. Hence the dimension of $\mathcal{H} \otimes \mathbb{Q}$ over $\mathbb{Q}_p$ and that of $\mathcal{H}/p\mathcal{H}$ over $\mathbb{F}_p$ must be coincide. Therefore, the leading coefficient of $A_L(t)$ is not divisible by any p. Since the characteristic polynomial of the t-action on each $\mathbb{Q}_p^{\deg(A_L(t))}$ is $A_L(t)$, its p-adic entropy is given by $h_p = \log \mathrm{M}_p(A_L(t)) = 0$.

The pontryagin dual of $\prod_p \mathbb{Q}_p$ is isomorphic to $\prod_p \mathbb{Q}_p$. We can replace the adèle ring $\mathbb{A}_{\mathbb{Q}}$ by $\prod_p \mathbb{Q}_p$ in the argument of [LW88, Lemmas 4.3-4.5] with use of results of [Bow71] and [Wal82]. Since the finite set P of prime numbers with contribution is empty, the topological entropy h of the t-action on the dual of $\mathcal{H}_{\mathbb{Q}}$ is given by $h = 0$.

Acknowledgment. I would like to express my sincere gratitude to Mikio Furuta and Yuichiro Taguchi for their precise questions at the thesis defense. I am grateful to Teruhisa Kadokami, Yasushi Mizusawa, Takayuki Morisawa, Masanori Morishita, Hirofumi Niibo, Yuji Terashima, and people at the coffee time of Komaba for helpful comments. I would like to thank Tomoki Mihara for precious guidance to p-adic analysis. I am also grateful to Dohyeong Kim for inviting me to the beautiful environment of POSTECH IBS-CBP, Noburu Ito and Kouki Taniyama for serving an important opportunity to give a talk, and my family for essential support. I was partially supported by Grant-in-Aid for JSPS Fellows (25-2241).

References

[AKM65] R. L. Adler, A. G. Konheim, and M. H. McAndrew. Topological entropy. *Trans. Amer. Math. Soc.*, 114:309–319, 1965.

[BD99] Amnon Besser and Christopher Deninger. p-adic Mahler measures. *J. Reine Angew. Math.*, 517:19–50, 1999.

[Bow71] Rufus Bowen. Entropy for group endomorphisms and homogeneous spaces. *Trans. Amer. Math. Soc.*, 153:401–414, 1971.

[BV13] Nicolas Bergeron and Akshay Venkatesh. The asymptotic growth of torsion homology for arithmetic groups. *J. Inst. Math. Jussieu*, 12(2):391–447, 2013.

[Den09] Christopher Deninger. p-adic entropy and a p-adic Fuglede-Kadison determinant. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 423–442. Birkhäuser Boston, Inc., Boston, MA, 2009.

[DGB14] Dikran Dikranjan and Anna Giordano Bruno. The bridge theorem for totally disconnected LCA groups. *Topology Appl.*, 169:21–32, 2014.

[EW99] Graham Everest and Thomas Ward. *Heights of polynomials and entropy in algebraic dynamics*. Universitext. Springer-Verlag London, Ltd., London, 1999.

[Fox56] Ralph H. Fox. Free differential calculus. III. Subgroups. *Ann. of Math. (2)*, 64:407–419, 1956.

[GAS91] Francisco González-Acuña and Hamish Short. Cyclic branched coverings of knots and homology spheres. *Rev. Mat. Univ. Complut. Madrid*, 4(1):97–120, 1991.

[GBV15] Anna Giordano Bruno and Simone Virili. Algebraic Yuzvinski formula. *J. Algebra*, 423:114–147, 2015.

[Hil12] Jonathan Hillman. *Algebraic invariants of links*, volume 52 of *Series on Knots and Everything*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition, 2012.

[HMM06] Jonathan Hillman, Daniel Matei, and Masanori Morishita. Pro-p link groups and p-homology groups. In *Primes and knots*, volume 416 of *Contemp. Math.*, pages 121–136. Amer. Math. Soc., Providence, RI, 2006.

[Iwa59] Kenkichi Iwasawa. On Γ-extensions of algebraic number fields. *Bull. Amer. Math. Soc.*, 65:183–226, 1959.

[KM08] Teruhisa Kadokami and Yasushi Mizusawa. Iwasawa type formula for covers of a link in a rational homology sphere. *J. Knot Theory Ramifications*, 17(10):1199–1221, 2008.

[KM13] Teruhisa Kadokami and Yasushi Mizusawa. On the Iwasawa invariants of a link in the 3-sphere. *Kyushu J. Math.*, 67(1):215–226, 2013.

[Lal03] Matilde N. Lalín. Some examples of mahler measures as multiple polylogarithms. *J. Number Theory*, 103(1):85–108, 2003.

[Le14] Thang Le. Homology torsion growth and Mahler measure. *Comment. Math. Helv.*, 89(3):719–757, 2014.

[LW88] D. A. Lind and T. Ward. Automorphisms of solenoids and p-adic entropy. *Ergodic Theory Dynam. Systems*, 8(3):411–419, 1988.

[Mih12] Tomoki Mihara. Singular homology of non-archimedean analytic spaces and integration along cycles. preprint. arXiv:1211.1422v1, 2012.

[MM82] John P. Mayberry and Kunio Murasugi. Torsion-groups of abelian coverings of links. *Trans. Amer. Math. Soc.*, 271(1):143–173, 1982.

[Nog07] Akio Noguchi. Zeros of the Alexander polynomial of knot. *Osaka J. Math.*, 44(3):567–577, 2007.

[Por04] Joan Porti. Mayberry-Murasugi's formula for links in homology 3-spheres. *Proc. Amer. Math. Soc.*, 132(11):3423–3431 (electronic), 2004.

[Ril90] Robert Riley. Growth of order of homology of cyclic branched covers of knots. *Bull. London Math. Soc.*, 22(3):287–297, 1990.

[Rol76] Dale Rolfsen. *Knots and links*. Publish or Perish, Inc., Berkeley, Calif., 1976. Mathematics Lecture Series, No. 7.

[Sak81] Makoto Sakuma. On the polynomials of periodic links. *Math. Ann.*, 257(4):487–494, 1981.

[Shn38] Lev Genrikhovich Shnirel'man(Schnirelmann). Sur les fonctions dans les corps normés et algébriquement fermés. *Izv. Akad. Nauk SSSR Ser. Mat.*, 2(5-6):487–498 (Russian), 1938.

[Smy81] C. J. Smyth. On measures of polynomials in several variables. *Bull. Austral. Math. Soc.*, 23(1):49–63, 1981.

[SW02] Daniel S. Silver and Susan G. Williams. Mahler measure, links and homology growth. *Topology*, 41(5):979–991, 2002.

[Uek16a] Jun Ueki. On the Iwasawa invariants for links and Kida's formula. preprint. arXiv:1605.09036, 5 2016.

[Uek16b] Jun Ueki. On the Iwasawa μ-invariants of branched $\mathbb{Z}_p$-covers. *Proc. Japan Acad. Ser. A Math. Sci.*, 92(6):67–72, 2016.

[Uek17] Jun Ueki. The profinite completions of knot groups determine the alexander polynomials. preprint. arXiv:1702.03836, 2 2017.

[Wal82] Peter Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.

[Was97] Lawrence C. Washington. *Introduction to cyclotomic fields*, volume 83 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1997.

[Web79] Claude Weber. Sur une formule de R. H. Fox concernant l'homologie des revêtements cycliques. *Enseign. Math. (2)*, 25(3-4):261–272 (1980), 1979.

Jun Ueki
Graduate School of Mathematical Sciences, The University of Tokyo,
3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan
E-mail: uekijun46@gmail.com